PCT/EP2004/007889

# IAP20 Rec'd FETAPTO 17 JAN 2006

# A cleansing composition based on oily substances

The present invention relates to a novel cleansing composition based on oily substances, particularly for cleansing skin and/or hair, which has excellent dermatological properties.

Skin and hair can be cleansed in various ways, even with water alone. However, the use of water alone may cause swelling of the horny layer and depletion of the water-soluble constituents of the NMF (Natural Moisturizing Factor), that is, the group of water-soluble substances which are responsible for correct skin hydration. Moreover, water alone cannot remove dirt of an oily nature or any residues of a lipid nature.

For cleansing skin and hair, it is therefore usual to use either formulations based on water and surfactants, that is, molecules which, by virtue of their amphiphilic structure, are capable of removing lipid substances and keep them in aqueous solution or suspension, or formulations based on oily substances which make use of the "similia similibus solvuntur" principle. By virtue of their oily nature, these substances are in fact capable of removing sebum and hydrophobic dirt without, however, removing the watersoluble hydrating components (NMF) of the skin. particularly desirable when the skin or hair are irritated by environmental damaged conditions (for example, atmospheric conditions pollution) or orphysiological conditions (for example, ageing or sensitive skin).

There are many oily substances which can be used for this purpose and they may be either natural or synthetic. Amongst these are substances which are fluid at ambient temperature (oils), pasty substances (butters, fats), and

solid substances (waxes). From the chemical point of view, these substances can be classified as:

- hydrocarbons (paraffinic or naphthenic) such as mineral oil, vaseline, paraffin, ozocerite, ceresin;
- terpenic hydrocarbons such as squalane, squalene, pristane, hexaisoprene, polyisobutylene;
- siloxanic polymers such as silicones and other heterolipids, including propoxylated alcohols;
- triglyceric esters such as most vegetable and some animal oils and fats (groundnut oil, karite butter, safflower oil, coconut butter, palm butter, soya oil, sunflower oil, olive oil, rice-germ oil, sweet almond oil, sesame oil, cocoa butter, wheat-germ oil, grape-seed oil, etc.); lecithin, which is a phosphatidyl cholinic glyceride, falls within this group;
- non-triglyceric esters such as lanolin, beeswax, spermaceti, carnauba wax, candelilla wax, the uropygial fat of aquatic birds, and a great many synthetic esters (isopropyl myristate, which is most often used, may be mentioned by way of example);
- fatty acids produced by the hydrolysis of vegetable fats (stearic acid is mainly used) of or lanolin (lanolic acids);
- fatty alcohols (for example, oleyl alcohol, cetyl alcohol) sterolic alcohols (for example, cholesterol), triterpenic alcohols (for example, lanosterol); many of these alcohols form part of the non-saponifiable fraction of vegetable lipids; the group also includes synthetic alcohols such as octyl dodecanol;

- essential oils, for example, essential oil of rosemary, lavender, angelica, artemisia, valerian, basil, bergamot, citronella, lemon, myrrh, patchouli, rose, or cinnamon.

However, cleansing compositions which are based on oily substances pose considerable formulation problems which are due both to poor solubility in water and to low foaming capacity. These problems have been addressed in the prior art by the use of various surfactants capable of increasing the dispersibility of the oily substances in water, improving the rinsability of the composition, and/or increasing foam formation.

For example, patent application DE-A 29 43 202 describes the use of monoalkylamines or dialkylamines, monoalkanolamines or dialkanolamines and/or salts of alkyl/alkanolamines of fatty alcohol sulphates as surfactants.

US Patent No. 4,371,548 describes the use of a  $C_8$ - $C_{18}$  sulphated fatty alcohol, optionally ethoxylated, neutralized by one or more amines, in a mixture with a  $C_8$ - $C_{10}$  ethoxylated sulphated fatty alcohol of ammonium.

US Patent No. 5,653,988 describes the use of a mixture constituted by a fatty alcohol, optionally ethoxylated, neutralized with various amines (preferably MIPA), an ethoxylated fatty alcohol (preferably Laureth-4) and a mono/diethanol amide (preferably Cocamide DEA).

US Patent No. 6,132,738 also describes the use of TIPA Laureth Sulphate.

However, a further problem to be addressed in the formulation of cleansing compositions for skin and/or hair is that of the aggressiveness of the surfactant. In the

cosmetics field, there is in fact a tendency to avoid the use of ingredients, particularly surfactants, which are irritating to skin and hair. This is even more desirable for compositions which are based on oily substances, the beneficial effects of which would be cancelled out by the presence of an aggressive surfactant in the composition. For this reason, the current tendency in the cosmetics field is to avoid the use of aggressive surfactants such as, for example, cocamide DEA - because of the possible formation of nitrosamines - and sulphated fatty alcohols, because of their aggressiveness and the dioxane content of the ethoxylated derivatives.

The present inventors have now found that the use of acylates of amino-acids, proteins and/or peptides surfactants in the formulation of cleansing compositions based on oily substances, particularly for cleansing skin and/or hair, enables compositions having excellent dermatological properties to be produced. These compositions have in fact been found to be extremely gentle towards skin and/or hair.

Simply adding these surfactants to a conventional oily formulation can reduce its aggressiveness and increase its tolerability. These molecules may also have a corrosion-inhibiting effect on domestic pipes.

A subject of the present invention is therefore a cleansing composition, particularly for cleansing skin and/or hair, comprising one or more oily substances at a total concentration within the range of between 10% and 90% by weight relative to the composition, preferably between 30% and 70%, and a surfactant selected from N-acylates of aminoacids, proteins, and peptides, and mixtures thereof, at a total concentration within the range of between 2% and 80%

by weight relative to the composition, said N-acylates of amino-acids, proteins, and peptides having formula (I):

$$\begin{array}{c}
O \\
|| \\
R-C \longrightarrow \\
\text{amino-acid residue} \longrightarrow \\
\text{m}
\end{array}$$
OB
(I)

wherein:

R is a saturated or unsaturated, linear or branched radical with from 3 to 30 carbon atoms, and the acyl group R-CO- is bound by an amide bond to the amine-nitrogen of the amino-acid residue,

m is a whole number  $\geq$  1, in which, when m>1, the repeated amino-acid residues may be identical to or different from one another,

B is hydrogen or a cation of an inorganic or organic base,

the composition being substantially anhydrous or having a water content no greater than 10% by weight.

In formula (I) above, the expression "amino-acid residue" includes both residues of non-cyclic amino-acids and residues of cyclic amino-acids. An amino-acid residue can be represented schematically by formula (II):

in which R' is the side chain of the amino-acid.

In the N-acylates of formula (I) which are used in the present invention, the acyl group R-CO- is bound to the amine-nitrogen of the amino-acid residue and the -OB group is bound to the carboxylic carbon of the amino-acid residue.

The N-acylates which are used in the present invention may be in acid form (when B = H) or may be neutralized (when B is a cation of an inorganic or organic base).

When B is a cation of an organic base, the organic base is preferably selected from MIPA (mono-isopropanol amine), TIPA (tri-isopropanol amine), Tris Amino (2-amino-2hydroxymethyl-1,3-propandiol), AEPD (2-amino-2-ethylpropandiol), (aminomethyl propandiol) and AMPD AMP (aminomethyl propanol). Even more preferably, the organic base is AMP.

When B is a cation of an inorganic base, the inorganic base is preferably selected from KOH, NaOH and  $NH_3$ .

When, in formula (I), m is 1 (m = 1), the surfactant is an N-acylate of an amino-acid. In this case, the amino-acid residue is preferably selected from the group consisting of the amino-acid residues of alanine, arginine, aspartic acid, asparagine, glutamic acid, glutamine, glycine, histidine, hydroxyproline, leucine, isoleucine, lysine, methionine, phenylalanine, proline, serine, threonine, tyrosine, valine, cystine, cysteine, sarcosine and pyrrolidone carboxylic acid.

When, in formula (I), m is greater than 1 (m > 1), the surfactant is an N-acylate of a peptide or a protein. The peptides/proteins which are most preferred for this purpose are of vegetable origin (for example, rice, wheat, oats,

maize, etc.) and are produced by the hydrolysis of the protein fraction of the plants used.

If other carboxylic groups are present on the side chains of the amino-acid residues, they may be in acid or salified form.

Surfactants which are particularly preferred for use in the composition of the invention are those in which the R-CO-acyl group is oleoyl or cocoyl. Most preferred amongst these are acid N-cocoyl sarcosinate and acid N-oleoyl sarcosinate, which can solubilize the oily substances contained in the composition, up to high concentrations, without the need for heating.

The surfactants which are used in the composition of the invention are synthesized by Schotten-Baumann reaction, starting with the chloride of the acid to be produced and with the desired amino-acid or peptide, or protein, in a basic NaOH or KOH environment. Upon completion of the reaction, acidification is then performed so as to obtain two phases, one constituted by the amino-acid or peptide or protein acylate in acid form and the other by water and sodium or potassium chloride (secondary reaction product). The acylate thus obtained can be used as such for the formulation of the cleansing composition neutralized beforehand with a base as described above. Alternatively, the acylate in acid form may be added directly to the formulation and may then be neutralized subsequently in the formulation.

The use of the surfactants described above has been found to be extremely effective. These surfactants can in fact solubilize oily substances up to high concentrations (up to 90% by weight relative to the formulation), even without the

need for heating, in the case of acid cocoyl and oleoyl sarcosinate. They also ensure optimal foaming and dermatological properties.

In some cases, to improve the clearness of the composition, it may be desirable to add an ethoxylated non-ionic surfactant, preferably an ethoxylated fatty alcohol.

In the cleansing composition of the invention, the above described N-acylates of amino-acids, peptides, or proteins can therefore be used individually or in mixtures. Their total concentration in the composition is within the range of from 2% to 80% by weight, preferably from 10% to 40% by weight, relative to the composition.

The cleansing composition of the invention may also optionally include further additives or active ingredients which are conventional for cosmetic compositions such as, for example, viscosifiers, colours, perfumes, antioxidants, and preservatives, the selection and correct use of which fall within the capabilities of a person skilled in the art and are therefore not described herein since they do not form a specific subject of the invention.

The cleansing compositions of the invention are prepared in the following manner:

- mixing together the oily substances to be used;
- dissolving the appropriate fragrance (it is important that it does not give rise to cloudiness or incompatibility); if an ethoxylated non-ionic surfactant is used, dissolving the fragrance therein;

- dissolving the desired surfactant/s and homogenizing to clearness;
- optionally neutralizing the surfactant/s in acid form by the addition of a neutralizing base upon completion of the formulation.

Some examples of oily cleansing compositions according to the invention are given below. These examples are provided for illustrative purposes and are not intended in any way to limit the scope of the invention as defined in the appended claims.

## **EXAMPLES**

### Example 1: bath oil

Acid oleoyl sarcosinate	26%
Perfume	1%
Soya oil	to 100%
Antioxidant	qds

## Example 2: shower oil

Acid cocoyl sarcosinate	10%
Perfume	5%
Isopropyl palmitate	15%
Paraffin oil	to 100%

## Example 3: oily face wash

Laureth-4	20%
Acid lauroyl glutamate	20%
Olive oil	48%
Isopropyl palmitate	10%

Perfume	2%
Example 4: shower oil	•
Cocoyl sarcosinate neutralized with AMP	37%
Oleoyl sarcosinate neutralized with AMP	30%
Wheat-germ oil	3%
Vaseline oil FU (Farmacopea Ufficiale -	
Italian Official Pharmacopoeia)	25%
Essence	5%
Antioxidants	ads

WO 2005/013927

#### CLAIMS

1. A cleansing composition, particularly for cleansing skin and/or hair, comprising one or more oily substances at a total concentration within the range of between 10% and 90% by weight relative to the composition and a surfactant selected from N-acylates of amino-acids, proteins and peptides, and mixtures thereof, at a total concentration within the range of between 2% and 80% by weight relative to the composition, the N-acylates of amino-acids, proteins and peptides having formula (I):

$$\begin{array}{c}
O \\
|| \\
R-C \longrightarrow \text{amino-acid residue} \longrightarrow OB \\
\end{array}$$
(I)

wherein:

R is a saturated or unsaturated, linear or branched radical with from 3 to 30 carbon atoms, and the acyl group R-CO- is bound by an amide bond to the amine-nitrogen of the amino-acid residue,

m is a whole number  $\geq$  1, in which, when m > 1, the repeated amino-acid residues may be identical to or different from one another,

B is hydrogen or a cation of an inorganic or organic base,

the composition being substantially anhydrous or having a water content no greater than 10% by weight relative to the composition.

WO 2005/013927 PCT/EP2004/007889

- 2. A cleansing composition according to Claim 1, wherein the oily substance is at a total concentration within the range of between 30% and 70% by weight, relative to the composition.
- 3. A cleansing composition according to Claim 1 or Claim 2, wherein the surfactant is at a concentration within the range of between 10% and 40% by weight, relative to the composition.
- 4. A cleansing composition according to any one of Claims 1 to 3, wherein B is a cation of an organic base selected from MIPA (mono-isopropanol amine), TIPA (tri-isopropanol amine), tris amino (2-amino-2-hydroxymethyl-1,3-propandiol), AEPD (2-amino-2-ethyl-1,3-propandiol), AMPD (aminomethyl propandiol) and AMP (aminomethyl propanol) or a cation of an inorganic base selected from KOH, NaOH and NH<sub>3</sub>.
- 5. A cleansing composition according to any one of Claims 1 to 4, wherein the surfactant is selected from the N-acylates of alanine, arginine, aspartic acid, asparagine, glutamic acid, glutamine, glycine, histidine, hydroxyproline, leucine, isoleucine, lysine, methionine, phenylalanine, proline, serine, threonine, tyrosine, valine, cystine, cysteine, sarcosine, pyrrolidone carboxylic acid, and mixtures thereof.
- 6. A cleansing composition according to any one of Claims 1 to 4, wherein the surfactant is selected from N-acylates of proteins and peptides of vegetable origin, preferably from rice, wheat, oats, maize, or mixtures thereof.
- 7. A cleansing composition according to any one of Claims 1 to 4, wherein the surfactant is selected from acid N-cocoyl

sarcosinate, acid N-oleoyl sarcosinate, and mixtures thereof.

- 8. A cleansing composition according to any one of Claims 1 to 7, wherein the oily substance is selected from the group consisting of paraffinic hydrocarbons, naphthenic hydrocarbons, terpenic hydrocarbons, siloxanic polymers, triglyceric esters, non-triglyceric esters, fatty acids, fatty alcohols, essential oils, and mixtures thereof.
- 9. A cleansing composition according to any one of Claims 1 to 8, further comprising a further, non-ionic ethoxylated surfactant, preferably an ethoxylated fatty alcohol, or a mixture of non-ionic ethoxylated surfactants.
- 10. Use of N-acylates of amino-acids, proteins, and peptides as defined in any one of Claims 1, 4, 5, 6 and 7 as surfactants in cleansing compositions based on oily substances, particularly for cleansing skin and/or hair.

A. CLASSIFICATION OF SUBJECT MATTER IPC 7 A61K7/06 A61K7/48 A61K7/50

According to International Patent Classification (IPC) or to both national classification and IPC

#### **B. FIELDS SEARCHED**

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

EPO-Internal, WPI Data, PAJ, CHEM ABS Data

Category °	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to daim No.
Х	US 5 843 407 A (LEATHERBURY NEIL CAMPBELL ET AL) 1 December 1998 (1998-12-01)	1-10
	column 1, line 56 - column 2, line 11 column 5, line 6 - line 14 column 7, line 40 - column 8, line 13 column 11, line 31 - line 51 column 12, line 34 - line 38 column 16, line 16 - line 18 examples VII,VIII	
Y	WO 98/09611 A (SEPPIC SA) 12 March 1998 (1998-03-12) page 1, line 4 - line 32 page 3, line 1 - line 26 page 4, line 15 - line 29 examples	1-10
	-/	

X Further documents are listed in the continuation of box C.	Patent family members are listed in annex.
Special categories of cited documents:  'A' document defining the general state of the art which is not considered to be of particular relevance  'E' earlier document but published on or after the international filing date  'L' document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)  'O' document referring to an oral disclosure, use, exhibition or other means  'P' document published prior to the international filing date but later than the priority date claimed	<ul> <li>"T" later document published after the International filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention</li> <li>"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone</li> <li>"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.</li> <li>"&amp;" document member of the same patent family</li> </ul>
Date of the actual completion of the international search  24 November 2004	Date of mailing of the international search report  15/12/2004
Name and malling address of the ISA  European Patent Office, P.B. 5818 Patentlaan 2  NL - 2280 HV Rijswijk  Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,  Fax: (+31-70) 340-3016	Authorized officer Ruckebusch, V

Category °	ation) DOCUMENTS CONSIDERED TO BE RELEVANT	I Delivertie de la
Calegory *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	WO 03/039496 A (ARIOTTO ANGELO; GUALA FABRIZIO (IT); VILLA GIOVANNI (IT); MERLO ELISA) 15 May 2003 (2003-05-15) page 2, line 5 - line 25 page 4, line 15 - page 5, line 8 page 7, line 32 - page 8, line 30 examples	1-10
Υ	WO 98/29094 A (UCHIYAMA HIROTAKA; HU FANG (JP); HARADA KIROH (JP); SAKO TAKASHI (JP)) 9 July 1998 (1998-07-09) page 2, line 15 - line 36 page 3, line 14 - page 5, line 22 examples	1-10
Y	PATENT ABSTRACTS OF JAPAN vol. 2000, no. 04, 31 August 2000 (2000-08-31) & JP 2000 007537 A (ASAHI CHEM IND CO LTD), 11 January 2000 (2000-01-11) abstract	1-10
Y	PATENT ABSTRACTS OF JAPAN vol. 0134, no. 02 (C-633), 6 September 1989 (1989-09-06) & JP 1 146816 A (LION CORP), 8 June 1989 (1989-06-08) abstract	1-10
Ρ,Χ	US 2003/162687 A1 (YAMAGUCHI YOSHINORI ET AL) 28 August 2003 (2003-08-28) page 1, paragraph 5 - paragraph 7 page 1, paragraph 8 - page 2, paragraph 9 page 2, paragraph 16 examples	1-10

International Application No

# /EP2004/007889

	_					
Patent document cited in search report		Publication date		Patent family member(s)		Publication date
US 5843407	A	01-12-1998	AU CA CN CZ DE DE EP ES JP WO EP	697809 8014994 2173104 1135171 9601107 69426887 0725620 2155097 9503785 3338057 9511000 0950392	A A1 A ,B A3 D1 T2 A1 T3 T B2 A1	15-10-1998 08-05-1995 27-04-1995 06-11-1996 14-08-1996 19-04-2001 04-10-2001 14-08-1996 01-05-2001 15-04-1997 28-10-2002 27-04-1995 20-10-1999
WO 9809611	A	12-03-1998	FR EP WO JP	2753096 0925058 9809611 2000517335	A1	13-03-1998 30-06-1999 12-03-1998 26-12-2000
WO 03039496	A	15-05-2003	IT WO EP	T020011053 03039496 1446092	A1	07-05-2003 15-05-2003 18-08-2004
WO 9829094	A	09-07-1998	WO AU BR CA EP JP US	9829094 1296597 9612850 2274849 0948315 2000505099 6284230	A A1 A1 T	09-07-1998 31-07-1998 28-02-2001 09-07-1998 13-10-1999 25-04-2000 04-09-2001
JP 2000007537	A	11-01-2000	NONE			
JP 1146816	Α	08-06-1989	JP	2609644	B2	14-05-1997
US 2003162687	A1	28-08-2003	FR JP	2836153 2003306696		22-08-2003 31-10-2003

# This Page is Inserted by IFW Indexing and Scanning Operations and is not part of the Official Record

# **BEST AVAILABLE IMAGES**

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images include but are not limited to the items checked:
☐ BLACK BORDERS
☐ IMAGE CUT OFF AT TOP, BOTTOM OR SIDES
FADED TEXT OR DRAWING
BLURRED OR ILLEGIBLE TEXT OR DRAWING
☐ SKEWED/SLANTED IMAGES
☐ COLOR OR BLACK AND WHITE PHOTOGRAPHS
☐ GRAY SCALE DOCUMENTS
LINES OR MARKS ON ORIGINAL DOCUMENT
REFERENCE(S) OR EXHIBIT(S) SUBMITTED ARE POOR QUALITY

# IMAGES ARE BEST AVAILABLE COPY.

As rescanning these documents will not correct the image problems checked, please do not report these problems to the IFW Image Problem Mailbox.